

Predicting diabetic ketoacidosis severity score: proposal of a therapeutic strategy adapted to emergency department

Houda Ben Soltane, Marwa Talbi, Mariem Khrouf, Ahmed Guesmi, Zied Mezgar, Mehdi Methamem.

Emergency department, Farhat Hached University Hospital, Sousse, Tunisia.

Corresponding author: Houda Ben Soltane; email: bensoltanehouda@yahoo.fr

Abstract

Background: We investigated independent severity predictors of diabetic ketoacidosis and developed a prediction rule for emergency physicians to classify patients into severity groups.

Methods: This study was done in a university-affiliated medical center. Consecutive adult patients (>15 years old) visiting the emergency department (ED) between July 2016 and July 2018 were enrolled when they met the criteria of DKA. Hospitalization in an intensive care setting was the primary endpoint.

Results: We included 176 patients diagnosed with DKA in the ED. We found 7 independent severity predictors: Altered mental status, venous pH, bicarbonate level, blood glucose, Serum creatinine, H4 serum chloremia and effective serum osmolality at admission. After assigning weights to each predictor, we developed a predicting DKA severity score that stratifies patients into groups: low severity (score<3): treatment in a general ward or the ED; and high (score≥3): treatment in the ICU. The area under the curve for the rule was 0.863.

Conclusions: The score is a simple and rapid rule for predicting severity and classifying patients with DKA.

Keywords: Diabetic Ketoacidosis; Score; Outcomes; Emergency

INTRODUCTION

Diabetes is a public health problem and pathology burdened with significant morbidity and mortality due to its acute and chronic complications.

Diabetic ketoacidosis (DKA) is a serious, potentially lethal complication of diabetes and may be the initial mode of presentation for patients with new-onset

T1D. It results from an absolute or relative insulin deficit with the immediate consequence of hyperglycemia and ketosis, which is itself responsible for water depletion and hydro-electrolyte disorders [2]. Regardless of the observed decrease in the death rate, DKA remain a serious health problem, especially in developing and undeveloped countries [1].

A better understanding of the pathophysiology combined with more uniform diagnostic and therapeutic treatment has led to a marked decline in developed countries, where its mortality varies between 0% and 5% [3, 4, and 5]. This metabolic disorder constitutes one of the main reasons for admission to the emergency department (ED) [1]. Several series studied mortality predictors and classified them in scores [6, 7, 8, and 9].

This study aimed to identify the clinical and evolutionary profile of the population admitted to the ED with a diagnosis of DKA, and use multivariate logistic regression analysis to investigate independent severity predictors and develop a prediction rule useful for ED physicians, one that allows them to classify patients with DKA into severity groups, and that allows them to make an appropriate management decision (treatment in the ICU, a general ward, or only in the ED).

METHODS

Study design, setting, population, and selection of participants: This study was done in a university-affiliated medical center in Tunisia with an ED staffed with board-certified emergency physicians. Consecutive adult patients (>15 years old) visiting the ED between July 2016 and July 2018 were enrolled when they met the following criteria [3]: DKA was defined as plasma glucose >2.5g/L, a high anion gap metabolic acidosis (serum HCO₃ <15 mmol/L, and pH >7.3), and positive urine ketones or serum ketones. Any other mode of decompensation: simple hyperglycemia, simple ketosis, hyperosmolar, a combination of hyperglycemia-

hyperosmolar state (osmolality > 320 mOsm/L) or a mixed syndrome ketoacetic and hyperosmolar was excluded. The effective serum osmolality was calculated with the formula: $2[\text{measured Na}^+ (\text{mEq/L})] + [\text{glucose} (\text{mg/dL})]/18$.

Data collection: Patients were prospectively selected in the ED. Insufficient information was retrospectively collected by checking medical records after the patients had been discharged from the hospital. The processing of the files was carried out based on a pre-established sheet. The data has been grouped into five themes: General data (age, sex, background, and the evolutionary profile of diabetes); clinical data; biological data at admission, at H4, at H8 and H12; and the analysis of triggering factors (therapeutic errors, infectious states, etc.). Therapeutic management and progressive characteristics (length of stay, mortality, and complications) were noted. We collected the quantity of insulin administered during the first 24 hours, the methods of its administration, as well as the other adjuvant therapies: rehydration, and antibiotic treatment.

Statistical analysis: All analyses were done using SPSS 25.0 for Windows. Descriptive statistics (mean and standard deviation [SD] for continuous variables and frequencies and proportions for categorical variables) were calculated.

Comparisons between the two groups were made using either an independent-samples t-test (assuming normal distribution) for the continuous variables.

Either a χ^2 test was used for categorical variables.

Odds ratios (ORs), along with the corresponding 95% confidence intervals (CIs), were also computed as appropriate. The area under the receiver operating characteristic (ROC) curves was used to compare a model's specifications along with its sensitivity and specificity. The results of the multivariate stepwise (forward) logistic regression analysis were then used to develop a prediction rule. Weights were assigned to each predictor according to their predicting β values of multiple logistic regression analysis. A Predicting DKA severity score was calculated for each patient. The scores on each different weight were used to determine their respective cutoff points for risk stratification by the ROC curve with the highest sum of sensitivity and specificity. For all analyses, a result was considered statistically significant at the $P < .05$ level of significance.

RESULTS

A total of 176 cases of DKA were included from 2016 to 2018. The mean age was 35.3 +/- 18.6 years, 32.4% were aged from 20 to 30 years, and 77 patients were male (sex ratio = 0.77). Diabetes was known in 150 patients (85.2%) of whom 60.8% were type 1 diabetics. 14.8% of patients were admitted for a primary diagnosis of DKA. The most common symptoms at presentation were abdominal pain (41.5%), and lethargy (13.1%). Other signs and symptoms at presentation were fever (9.1%), polyuria and polydipsia (8%), Kussmaul breathing (5.7%), and chest pain (1.7%). The most frequent decompensation factors were infection in 36.9%

and discontinuation of treatment in 23.9% of cases.

The average blood glucose was 28 +/- 7.6 mmol/L, the mean venous pH was 7.14 +/- 0.12, and blood HCO₃⁻ was 7.9 +/- 3.98 mmol / L. The main hydration solution was a 0.9% saline solution. The mean serum amount infused was 3.90 +/- 2.02 L / 24h. The mean dose of intravenous insulin infused in the first 24 hours was 43.94 +/- 12.74 IU / 24h. The mean time to switch to subcutaneous insulin therapy was 24.8 +/- 16.4 hours. Only five patients with hypoglycaemia were noted during treatment (2.8%). The length of stay in the ED was 16.1 +/- 13.5 hours. 19.7% of patients were discharged out of the ED, after clinical and biological recovery as well as the control of the cause of decompensation. In contrast, 121 patients (68.8%) were transferred to an endocrinology ward, and 13 (7.4%) were admitted to the intensive care unit. The average total hospital length of stay was 6.41 +/- 6 days. Five cases of death (2.8%) were described in our study. Then patients were prospectively assigned to 2 groups as follows: 1) the first group consisted of 13 patients who were admitted to the intensive care unit; 2) the second group consisted of 121 patients who were admitted to the General ward (endocrinology). They were compared to detect patients requiring intensive care upon admission to the emergency room (Table 1).

A multiple logistic regression analysis with stepwise variable selection using backward selection was performed and odds ratios with 95% confidence intervals were calculated for each predictor.

Table 1: Comparison between the ICU group and the Endocrinology Ward Group

Measures	Patients in ICU	Patients in Ward	P Values
Number	13	121	-
Age	37.2±19.2	35.1±18.6	0.694
DM duration	7.5±5.8	7±6.8	0.773
Males	7.8%	62.3%	0.106
PR	107.5±20.5	109.2±21.1	0.784
SBP	12.7±1.7	12.1±2.3	0.395
DBP	6.8±1.4	4±1.4	0.693
RR	27.8±3.8	25.9±6.8	0.297
GCS	14.2±2	14.8±0.7	0.002
Blood glucose	34.3 ± 8.5	27.6 ±7.4	0.002
Urea	11.7 ± 11	7.2± 4.8	0.02
Serum creatinine	166±196	94.7±89.4	0.025
Na+	132.3±6.4	131.6±4.8	0.62
K+	4.7±0.9	4.5±0.9	0.3
pH H0	7±0.10	7.20±0.10	p<0.001
HCO3- H0	4.5±3.4	8.2± 3.9	0.001
Osmolarity	298.9±14.4	290.7±10.7	0.01
pH H4	7.20 ± 0.10	7.30 ± 0.10	0.001
HCO3- H4	8.1 ± 3.5	12.6± 4.7	0.001
Na+ H4	134.1 ± 6.4	134±4.6	0.95
K+ H4	4 ± 0.8	3.7± 0.8	0.218
Cl H4	110.1 ± 6.4	103.7± 3.7	0.005
pH H8	7.20± 0.10	7.30 ±0.10	<0.001
HCO3- H8	10.3 ±4.3	14.5 ±4.3	0.002
pH H12	7.20 ± 0.10	7.30 ± 0.10	0.003
HCO3- H12	11.2 ± 5.1	15.2 ± 4.2	0.007
Na+ H12	133.8 ± 6.4	135± 4.2	0.396
K+ H12	3.8± 0.9	3.7 ±0.7	0.466
Cl H12	111 ± 5.4	105 ± 6.4	0.007
Mortality rate	4	1	<0.001

DM= diabetes mellitus; ICU= intensive care unit; PR=pulse rate; RR= respiratory rate; SBP= systolic blood pressure; DBP=diastolic blood pressure;

The results were used to develop a clinical and biological severity score, to detect patients who need admission in ICU.

After this analysis, 7 independent severity predictors were retained: Altered mental status, venous pH, bicarbonate level, blood glucose, Serum creatinine, H4 serum chloremia, and effective serum osmolality at admission. A weight was assigned to each element of this score according to its predictive β value (table 2).

Table 2: Predicting DKA Severity Score

Variable	weights
Glasgow Coma Scale (GCS) < 15	1
Venous pH < 7.14	2
Serum HCO3- < 7.7 m moles/l	1
Blood glucose > 28,1 m moles/l	1
Serum creatinine > 100	1
Chloremia at H4 > 104	1
Serum Osmolarity > 291m osm/l	1

Low severity: score<3; High severity: score≥3

This score will allow ED physicians to classify patients according to their clinical and biological gravity and therefore to better manage them in the most appropriate environment. Finally, it was calculated for each patient in our population. The mean score was 5.5 ± 1.4 for those who were admitted to the intensive care unit, and 2.8 ± 1.9 for the rest of the patients ($p = 0.0001$).

Our population can then be classified into two severity groups: low with a score of <3 and high with a score ≥ 3 [OR (95% CI), 10.7 (1.36-84.5)]. A Score greater than 3 is predictive of severity. It has a high predictive power (The area under the ROC curve is 0.863, with 100% sensitivity and 71.2% specificity) (Figure 1).




Figure 1: Presentation of the sensitivity and specificity of the severity score

DISCUSSION

In Tunisia, the prevalence of diabetics was passed from 9% in 2007 to 15% in 2017. The age of onset is getting younger and younger, which exposes people to more acute and chronic complications [10,11,12]. Acute complications of diabetes are a reason for frequent consultation and hospitalization in emergency departments [12,13]. Among these complications, is ketoacidosis (DKA) which is an acute and severe complication, given the importance of its morbidity and mortality [3,14]. Its annual incidence in the USA is estimated between 4.6 and 8 episodes/1000 diabetics with a constant increase [10,12,15,16]. This incidence is even greater in other countries, notably in England, Sweden, and the developing countries [15]. In a large study published in December 2015, admissions for DKA represented 1.1% of all ICU

admissions in Australia and New Ireland between 2000 and 2013 [17].

In prior studies, predictors of DKA mortality were studied and scored: A study carried out in Parakou Hospital ICU in Benin over a period of 10 months, showed that the occurrence of complications was associated with advanced age, low socioeconomic level, high serum osmolarity, and long consultation time. It objectified a mortality rate of 14%, having as predictive factors the advanced age and long period before consultation [6]. Another study done in Taiwan developed a mortality predictor score (PHD) finding 6 independent mortality predictors: Absent tachycardia, Hypotension, Anemia, Severe coma, Cancer history, and Infection [7]. The study by Stéphanie T. Chung (Jamaica 2006) concluded that mortality increased significantly with age and that the main mortality predictors were altered mental status on admission, co-morbidity, age, diabetes duration, and association of DKA with hyperosmolar syndrome [8]. For Stamatis P. et al, mortality predictors were co-morbidity, pH <7.0, a total dose of IV insulin injected in the first 12 hours > 50 IU, blood glucose > 16.7 mmol / L after 12 hours of treatment, altered mental status, and persistent fever after 24 hours [18]. Several series studied mortality predictors and ranked them as scores, but there were enough series that studied severity predictors and admission risk ICU.

We developed a novel decision rule to predict severity and manage ED patients with DKA. ED and ICU physicians can usefully evaluate 7 variables. Patients with a high score should be

deemed critically ill and sent to the ICU for advanced treatment such as aggressive fluid resuscitation, strict intravenous insulin control, detailed investigation and management of the precipitating factors, and careful prevention of possible treatment complications. For patients with low scores, a general ward admission or ED treatment may be sufficient, which would help preserve medical resources for patients in greater need.

Blood PH at admission, with a high β -value, was the strongest severity predictor. Altered mental status has been proposed as the only independent severity predictor, which is easier to quantify in clinical practice. Blood PH and effective serum osmolality are important for evaluating DKA; these two factors appear to be significant severity predictors in our study. Blood creatinine, which reflects the importance of water deficit, can be at the origin of functional renal failure [19]. It can also be falsely high due to the presence of ketone bodies. Chloremia at H4 is a variable that was kept in our score as a severity factor. Indeed, rehydration with isotonic saline serum promotes the appearance of hyperchloremic acidosis which should be avoided [3].

LIMITATIONS

This study has several limitations. First, some clinical presentations or records may not have been completely documented. Second, this was a single-center study. Third, the sample size might not be large enough to make conclusions with good statistical power. Additional studies with larger sample sizes are necessary. Fourth, although we have validated the prediction rule in

a prospective cohort, external validation in other populations is necessary.

CONCLUSION

Diabetic ketoacidosis occurs in young people treated with insulin therapy. Infection appears to be the most implicated factor in decompensation. A Score greater than 3 is predictive of severity, requiring care in an ICU with a mortality rate remaining low thanks to this score.

REFERENCES

1. WHO. Diabète. 2023. [<https://www.who.int/fr/news-room/fact-sheets/detail/diabetes>]
2. Kitabchi, A. E., Umpierrez, G. E., Miles, J. M., & Fisher, J. N. Hyperglycemic crises in adult patients with diabetes. *Diabetes care*. 2009; 32: 1335-1343.
3. Kitabchi, A. E., Murphy, M. B., Spencer, J., Matteri, R., & Karas, J. Is a priming dose of insulin necessary in a low-dose insulin protocol for the treatment of diabetic ketoacidosis? *Diabetes care*. 2008;. 31: 2081-2085.
4. American Diabetes Associatio. Hyperglycemic crises in diabetes. *Diabetes Care*. 2004;27: s94-s102.
5. Maldonado, M. R., Chong, E. R., Oehl, M. A., & Balasubramanyam, A. Economic impact of diabetic ketoacidosis in a multiethnic indigent population: analysis of costs based on the precipitating cause. *Diabetes care*. 2003;26: 1265-1269.
6. Tchaou, B. A., Gomina, M., Agbo, A. H. M., & Akpona, S. A. Complications aigues métaboliques du diabète sucre dans l'unité de réanimation de l'hôpital universitaire de parakou (benin). *European Scientific Journal*, 2014;10
7. Huang, C. C., Kuo, S. C., Chien, T. W., Lin, H. J., Guo, H. R., Chen, W. L., ... & Su, S. B. Predicting the hyperglycemic crisis death (PHD) score: a new decision rule for emergency and critical care. *The American journal of emergency medicine*. 2013; 31: 830-834.
8. Chung, S. T., Perue, G. G., Johnson, A., Younger, N., Hoo, C. S., Pascoe, R. W., & Boyne, M. S. Predictors of hyperglycaemic crises and their associated mortality in Jamaica. *Diabetes research and clinical practice*. 2006; 73:184-190.
9. Chen, H. F., Wang, C. Y., Lee, H. Y., See, T. T., Chen, M. H., Jiang, J. Y., ... & Li, C. Y. Short-term case fatality rate and associated factors among inpatients with diabetic ketoacidosis and hyperglycemic hyperosmolar state: a hospital-based analysis over 15 years. *Internal Medicine*. 2010;49: 729-737.
10. Van Ness-Otunnu, R., & Hack, J. B. Hyperglycemic crisis. *The Jour of Emerg Med*. 2013;45: 797-805.

11. Gosmanov, A. R., Gosmanova, E. O., & Kitabchi, A. E). *Hyperglycemic crises: diabetic ketoacidosis (DKA), and hyperglycemic hyperosmolar state (HHS)*. 2021. In: Feingold KR, Anawalt B, Blackman MR, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-. Available from: [https://www.ncbi.nlm.nih.gov/books/NBK279052/Zhang, P., Zhang, X., Brown, J., Vistisen, D., Sicree, R., Shaw, J., & Nichols, G. \(2010\). Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes research and clinical practice*, 87\(3\), 293-301.](https://www.ncbi.nlm.nih.gov/books/NBK279052/Zhang, P., Zhang, X., Brown, J., Vistisen, D., Sicree, R., Shaw, J., & Nichols, G. (2010). Global healthcare expenditure on diabetes for 2010 and 2030. Diabetes research and clinical practice, 87(3), 293-301.)
12. Mazer, M., & Chen, E. *Is subcutaneous administration of rapid-acting insulin as effective as intravenous insulin for treating diabetic ketoacidosis?* *Annals of Emergency Medicine*. 2009; 53: 259-263.
13. Foster, J. R., Morrison, G., & Fraser, D. D.. *Diabetic ketoacidosis-associated stroke in children and youth*. *Stroke Res Treat*. 2011; 2011: 219706..
14. Nyenwe, E. A., & Kitabchi, A. E. *Evidence-based management of hyperglycemic emergencies in diabetes mellitus*. *Diabetes research and clinical practice*. 2011; 94.; 340-351.
15. Abdulrahman, G. O., Amphlett, B., & Okosieme, O. E.. *Trends in hospital admissions with diabetic ketoacidosis in Wales, 1999–2010*. *Diabetes research and clinical practice*. 2013; 100, e7-e10.
16. Venkatesh, B., Pilcher, D., Prins, J., Bellomo, R., Morgan, T. J., & Bailey, M. *Incidence and outcome of adults with diabetic ketoacidosis admitted to ICUs in Australia and New Zealand*. *Critical Care*. 2015; 10:451.
17. Efstathiou, S. P., Tsiakou, A. G., Tsioulos, D. I., Zacharos, I. D., Mitromaras, A. G., Mastorantonakis, S. E., ... & Mountokalakis, T. D. (2002). *A mortality prediction model in diabetic ketoacidosis*. *Clin endocrinol*. 2002;57:595-601.
18. Chiasson, J. L., Aris-Jilwan, N., Bélanger, R., Bertrand, S., Beaugard, H., Ékoé, J. M., et al. *Diagnosis and treatment of diabetic ketoacidosis and the hyperglycemic hyperosmolar state*. *Canadian Medical Association Journal*. 2003; 168: 859-866.